

The Role of Micronutrients in Child Growth and Development

May 19, 2021

Shawn Baker

Lynne Ausman

Andrew Thorne-Lyman

Omar Dary



FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative

WELCOME TO THE ZOOM WEBINAR

The screenshot shows a Zoom webinar interface. The main video area is black. On the right, there is a 'Zoom Group Chat' panel. In the bottom toolbar, the 'Join Audio' button is circled in orange, with an arrow pointing to the text 'If you are unable to hear, connect your speakers by selecting "Join Audio"'. Another arrow points from the 'Chat' button in the toolbar to the chat panel, which contains the text 'Please use the chat box to introduce yourselves and share thoughts and comments by sending a message to "All panelists and attendees"'. The chat panel also shows a dropdown menu with 'To: All panelists and attendees' selected and a text input field labeled 'Type message here ...'.

If you are unable to hear, connect your speakers by selecting "Join Audio"

Please use the chat box to introduce yourselves and share thoughts and comments by sending a message to "All panelists and attendees"

To: All panelists and attendees
Type message here ...

For a better Zoom Webinar Experience, download the Zoom app to your device at <https://zoom.us/support/download>



Q&A AND CHAT

The screenshot displays a Zoom meeting interface. On the left, a large black area contains the text "Submit your questions for the panelists in the Q&A box" in orange. An orange arrow points from the "Q&A" icon in the bottom toolbar to a white Q&A box. The Q&A box has a title bar "Q&A" and a close button. Inside, it says "Welcome 🍌" and "Feel free to ask the host and panelists questions". Below this is a text input field labeled "Type your question here...". On the right, a "Zoom Group Chat" window is open. It contains the text "If you're having any technical difficulties, please send a message to 'All panelists' via the chat box and we will do our best to help resolve your issue" in blue. A blue arrow points from a blue circle around the "All panelists" selection in the chat's "To:" dropdown menu up to the text. The chat window also shows a "Type message here ..." field and a "Leave" button at the bottom.

Submit your questions for the panelists in the Q&A box

Q&A

Welcome 🍌

Feel free to ask the host and panelists questions

Type your question here...

Zoom Group Chat

If you're having any technical difficulties, please send a message to "All panelists" via the chat box and we will do our best to help resolve your issue


✓ All panelists
All panelists and attendees

To: All panelists

Type message here ...

Leave

FEED THE FUTURE INNOVATION LAB FOR NUTRITION



The screenshot shows a web browser window with the URL nutritioninnovationlab.org. The page features a dark background with white text. On the left, there is a main heading and two bulleted lists. On the right, there is a video player showing a group of people in a field. The video player has a play button and a progress bar. Below the video player, there are social media icons for Twitter, Facebook, and YouTube. The bottom of the browser window shows the Windows taskbar with various application icons and the system clock.

Welcome to the Innovation Lab | x +

nutritioninnovationlab.org

Apps Recherche - Norma... Lancaster Guidos Services Inc... Penguins MAPPPD Canvas Nutr 203 P.E with Joe | Wedn... Social bridging mat... Gardening Advice a... Science Modernist Pantry

We're building institutional capacity for analysis and policy formulation in developing countries

- » Human and Institutional Capacity Building

We're pursuing applied research that supports the goals of the U.S. Government's Feed the Future initiative

- » Nutrition and Agriculture Linkages
- » Program and Policy Processes
- » Biological Mechanisms

Feed The Future | Innovation Lab for Nutrition Info... Watch later Share

0:00 / 4:16 YouTube


17:26 29/10/2020

www.nutritioninnovationlab.org



FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative





FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative

U.S. GOVERNMENT PARTNERS





GLOBAL AND LOCAL PARTNERS



WEDNESDAY, MAY 19TH
9:00AM - 10:30AM (ET)

The Role of Micronutrients in Child Growth and Development



SHAWN BAKER
USAID



ANDREW THORNE-LYMAN
Johns Hopkins University



LYNNE AUSMAN
Tufts University



OMAR DARY
USAID

Iron and Vitamin A Levels in Pregnant Women and Birth Outcomes: Results from a Birth Cohort Study in Uganda

Lynne M. Ausman, Grace Namirembe, and Julieta Mezzano for the Nutrition Innovation Lab

UGANDA BIRTH COHORT STUDY

- Mother infant dyads (sub study of n=1244) were followed during pregnancy up to 12 months after birth
- Dyads were equally distributed in 16 subcounties in North and Southwest Uganda
- Questionnaires and anthropometric measures were taken at regular intervals in both mothers and infants.
- Blood samples for micronutrient concentrations (iron and vitamin A) and inflammatory markers were collected at birth and six months after birth for mother and infants, respectively
- This report presents the results of mother and infant characteristics at time of parturition




FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative

UGANDA STUDY DISTRICTS

South Western districts

Ibanda
Kabale
Kamwenge
Kanungu
Kasese
Kiryandongo
Kisoro
Masindi



Northern districts

Agago

Dokolo

Kole

Lira

Nebbi

Oyam

Pader



USAID
FROM THE AMERICAN PEOPLE



GERALD J. AND DOROTHY R.
Friedman School of
Nutrition Science and Policy

IRON AND VITAMIN A SUPPLEMENTATION IN UGANDA

- Standard practices mandate from Ministry of Health:
 - Iron and folic acid supplementation to pregnant women
 - Cooking oil fortified with Vitamin A in form of retinyl palmitate
 - Commercial wheat flour and maize fortified with iron and three other nutrients

<https://www.unbs.go.ug/news-highlights.php?news=104&read>

BRINDA CORRECTION FOR IRON AND VITAMIN A BIOMARKERS

- Inflammatory biomarkers CRP, AGP, and malaria alter the expression of ferritin, sTfR, and RBP biomarkers.
- Utility to predict amounts of absolute deficiency can be improved with a BRINDA adjustment of these biomarkers.

$$\text{Ferritin}_{\text{adjusted}} = \text{ferritin}_{\text{unadjusted}} - \beta_1 (\text{CRP}_{\text{obs}} - \text{CRP}_{\text{ref}}) - \beta_2 (\text{AGP}_{\text{obs}} - \text{AGP}_{\text{ref}}) - \beta_3 \text{malaria}$$

Namaste SM, Aaron GJ, Varadhan R, Peerson JM, Suchdev PS; BRINDA Working Group. Methodologic approach for the Biomarkers Reflecting Inflammation and Nutritional Determinants of Anemia (BRINDA) project. Am J Clin Nutr. 2017 Jul;106(Suppl 1):333S-347S



MATERNAL MICRONUTRIENT & INFLAMMATION BIOMARKERS AT PARTURITION

Variable	n	Mean	SD	Median	Min	Max
Ferritin (corrected) (µg/L)	1244	48.90	30.13	42.73	3.38	176.88
Ferritin (µg/L)	1244	67.14	42.64	56.95	3.59	327.34
sTfR (corrected) (mg/L)	1244	7.25	4.18	6.02	2.05	45.20
sTfR (mg/L)	1244	7.80	4.92	6.35	1.67	49.74
Hemoglobin at birth (g/dL)	1186	13.00	1.61	13.10	7.00	17.80
RBP (µmol/L)	1244	1.97	0.77	1.84	0.41	4.00
AGP (g/L)	1244	1.07	0.58	0.93	0.13	4.57
CRP (mg/L)	1244	7.73	17.87	1.90	0.01	277.11
Albumin (g/dL)	1209	3.78	0.80	3.85	1.16	6.86

IRON DEFICIENCY AND ANEMIA IN UGANDAN MOTHERS

Parameter of deficiency	Prevalence (%) (95% CI)	
	Uncorrected	Corrected *
Iron depleted stores (FER)	7.4% (6.0, 8.9)	12.3% (10.5, 14.3)*
Iron deficient erythropoiesis (sTfR)	26.7% (24.3, 29.2)	21.6% (19.4, 24.0)*
Depleted body iron Stores (BIS)	8.0% (6.6, 9.7)	10.5% (8.8, 12.3)#
Anemia (Hgb)	11.0% (9.2, 12.9)	13.8% (11.9, 15.9)^
Iron Deficiency Anemia (IDA)	---	4.5% (3.4, 5.8)

* Correction for inflammation using BRINDA Coefficient Regression method

BIS was calculated as the ratio of BRINDA-adjusted sTFR/FER

^ Anemia was adjusted for altitude



VITAMIN A DEFICIENCY IN UGANDAN MOTHERS

Parameter of deficiency [#]	Prevalence (%) (95% CI)
	Uncorrected*
Vitamin A deficiency (RBP <0.83 µmol/L)	3.1% (2.2, 4.3)
Vitamin A deficiency (RBP < 1.05 µmol/L)	8.5% (7.2, 9.9)
Moderate vitamin A deficiency (RBP <1.17 µmol/L)	12.2% (10.5, 14.2)

[#] Using three commonly used cut-offs for RBP.

^{*} BRINDA adjustment for inflammation not recommended in mothers or women of reproductive age





BIRTH OUTCOMES

Characteristics	n	Mean \pm SD
Birth weight (kg)	1244	3.26 \pm 0.49
Birth Length (cm)	1244	47.57 \pm 3.33
Weight-for-age Z score	1239	-0.12 \pm 1.00
Length-for-age Z score	1233	-1.00 \pm 1.73
Weight-for-length Z score	1006	0.54 \pm 1.74
BMI Z-score	1215	0.60 \pm 1.65
Gestational age (days)	1108	274.56 \pm 21.71

Prevalence Estimates (%)	
Underweight	2.4
Stunted	26.2
Wasted	7.8
Pre-term birth	19.5
Low birth weight	3.5



MATERNAL FERRITIN ($\mu\text{g/L}$) AND BIRTH OUTCOMES

	UNADJUSTED MODEL			ADJUSTED MODEL #			
	n	β estimate (SE)	p value	n	β estimate (SE)	p value	
Birth weight (kg)	1244	-0.03 (0.02)	0.153	1077	-0.03 (0.02)	0.126	
Weight-for-age Z score	1240	-0.05 (0.04)	0.196	1076	-0.05 (0.04)	0.175	
Length-for-age Z score	1240	-0.01 (0.07)	0.874	1072	-0.07 (0.04)	0.132	
Weight-for-length Z score	1020	-0.03 (0.08)	0.722	867	0.11 (0.06)	0.081	
		OR (95% CI)			OR (95% CI)		
Small for Gestational Age	1001	0.97 (0.76-1.24)	-	915	0.98 (0.72-1.34)	-	
Preterm	1403	1.11 (0.93-1.32)	-	1018	1.24 (0.92-1.65)	-	

* Log (ln) transformed and adjusted for inflammation using BRINDA

#Adjusted for maternal age, education, & height, wealth index, subcounty, infant sex, iron supplementation frequency



MATERNAL sTFR (mg/L)* AND BIRTH OUTCOMES

	UNADJUSTED MODEL			ADJUSTED MODEL #			
	n	β estimate (SE)	p value	n	β estimate (SE)	p value	
Birth weight (kg)	1244	0.03 (0.03)	0.371	1077	-0.02 (0.04)	0.662	
Weight-for-age Z score	1240	0.06 (0.07)	0.375	1076	-0.02 (0.07)	0.767	
Length-for-age Z score	1240	0.17 (0.11)	0.126	1072	0.10 (0.14)	0.484	
Weight-for-length Z score	1020	0.07 (0.13)	0.615	867	0.03 (0.13)	0.829	
	n	OR (95% CI)		n	OR (95% CI)		
Small for Gestational Age	1001	0.75 (0.49-1.15)	-	915	0.96 (0.59-1.59)	-	
Preterm	1403	0.86 (0.64-1.16)	-	1018	0.67 (0.48-0.94)*	-	

* Log (ln) transformed and adjusted for inflammation using BRINDA'

Adjusted for maternal age, education, & height, wealth index, subcounty, infant sex, iron supplementation frequency.



MATERNAL RBP ($\mu\text{mol/l}$) AND BIRTH OUTCOMES

	UNADJUSTED MODEL			ADJUSTED MODEL [#]		
	n	β (SE)	p value	n	β (SE)	p value
Birth weight (kg)	1244	0.01 (0.02)	0.485	1077	-0.01 (0.02)	0.576
Weight-for-age Z score	1240	0.01 (0.04)	0.759	1076	-0.03 (0.05)	0.523
Length-for-age Z score	1240	0.16 (0.07)	0.014	1072	0.12 (0.06)	0.030
Weight-for-length Z score	1020	-0.06 (0.07)	0.413	867	-0.02 (0.06)	0.709
	n	OR (95% CI)		n	OR (95% CI)	
Small for Gestational Age (days)	1001	1.19 (0.95-1.50)	-	915	1.25 (0.90-1.74)	-
Preterm	1403	0.83 (0.70-0.98)	-	1018	0.88 (0.71-1.10)	-

* Log (ln) transformed and not adjusted for inflammation.

Adjusted for maternal age, education, & height, wealth index, subcounty, infant sex, iron supplementation frequency.





FOOD SECURITY, VITAMIN A & IRON INTAKE


Characteristics	%
HFIAS - Food Secure	36.52
Vegetable solid fats	1.51
Vegetable liquid oils	22.00
Wheat consumption	6.22
Maize consumption	33.31
Vit. A rich food intake (plant)	39.10
Vit. A rich food intake (animal)	11.59
Iron rich food intake (non-heme)	93.62
Iron rich food intake (heme)	22.04
MDD-W - Consumption of ≥ 5 food categories	17.14
Iron suppl. during pregnancy	94.13

SUMMARY OF FINDINGS

- The BRINDA adjustment for FER shows an increase in iron deficiency from 7.4% to 12.3% in women
 - After adjustment, maternal FER levels are positively trend toward improved WLZ ($p=0.08$)
- The BRINDA adjustment for sTfR shows a decrease in iron-deficient erythropoiesis from 26.7% to 21.6% in women
 - After adjustment, sTfR is associated with lower odds of preterm births ($OR = 0.67$)
- Maternal Vitamin A deficiency as measured by RBP ranges from 3% to 8.5%.
- Maternal RBP was associated with increased LAZ of offspring ($p=0.03$).
- Deficiencies of both nutrients continue to exist in pregnant women in Uganda. Vitamin A intake from oils, and both animal and plant source foods appears to be low.
- Heme iron intake is low
- Only 17% of women consumed at least 5 food groups (MDD-W) suggesting the necessity for investment in programs to diversify diets.


Lifecycle Connections in Our Micronutrient Related Research in Nepal

Andrew Thorne-Lyman, ScD, MHS
Associate Scientist, Johns Hopkins Bloomberg School of Public Health





PATHWAYS FOR MICRONUTRIENTS



Study 1. The risk factors for anemia are consistent across three national surveys in Nepal

Monica M. Pasqualino, Andrew L. Thorne-Lyman, Swetha Manohar, Angela KC, Binod Shrestha, Ramesh Adhikari, Rolf D. Klemm, Keith P. West Jr

Current Developments in Nutrition (In Press)




FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative

PoSHAN SURVEYS

Study design

- Nationally representative sample obtained across the Mountains, Hills, and Tarai.
- Annual same-season community-based observational surveys conducted in the same 63 wards in 2013, 2014, 2016
- Data collected on individual, household, and community-level factors.



USAID
FROM THE AMERICAN PEOPLE



NARC
Nepal Agricultural Research Council



Tufts
UNIVERSITY

GERALD J. AND DOROTHY R.
Friedman School of
Nutrition Science and Policy



ANEMIA ASSESSMENT

Methods

- Random sample of children 6-59 months and mothers selected each year for anemia assessment using a Hb 201+ hemoglobinometer (HemoCue AB, Angelholm, Sweden).
- Anemia status defined based on WHO standard (hemoglobin concentration <11.0 g/dL) and adjusted for altitude.



Prevalence of anemia among children 6-59 months


	2013	2014	2016
National, n	809	796	865
% (95% CI)	63.4 (59.1, 67.5)	52.1 (47.0, 57.3)	59.5 (54.8, 64.1)
Mountains, n	148	135	133
% (95% CI)	62.3 (54.9, 69.7)	52.6 (42.9, 62.1)	58.6 (46.1, 70.2)
Hills, n	221	214	245
% (95% CI)	51.1 (43.2, 59.0)	38.8 (30.2, 48.1)	46.1 (38.1, 54.3)
Tarai, n	440	447	487
% (95% CI)	70.0 (64.3, 75.2)	58.4 (52.3, 64.2)	66.5 (60.5, 72.0)



FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative

Child-level risk factors for child anemia



Age reference level: 36-59 mo

Diarrhea: any in last 7 days

Meat intake: frequency in last 7 days; reference level: 8+ times

Deworming: last 12 months



USAID
FROM THE AMERICAN PEOPLE




GERALD J. AND DOROTHY R.
Friedman School of
Nutrition Science and Policy



FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative


Partially adjusted odds ratios, maternal-level risk factors



Education reference level: Higher secondary or more
Thin: MUAC <22.5 cm



USAID
FROM THE AMERICAN PEOPLE



NARC
Nepal Agricultural Research Council



Tufts
UNIVERSITY

GERALD J. AND DOROTHY R.
Friedman School of
Nutrition Science and Policy

STUDY 2. WHAT FACTORS ARE ASSOCIATED WITH LOW DIETARY DIVERSITY IN PREGNANCY?

- Cross-sectional study, 327 pregnant women
- Baglung district, urban municipality in the western hill region of Nepal
- Minimum Dietary Diversity for Women (MDD-W), ≥ 5 of 10 food groups past 24 hours
- 45% of women had low dietary diversity

PLOS ONE

RESEARCH ARTICLE

Factors associated with dietary diversity among pregnant women in the western hill region of Nepal: A community based cross-sectional study

Vintuna Shrestha^{1*}, Rajan Paudel², Dev Ram Sunuwar³, Andrew L. Thorne Lyman⁴, Swetha Manohar^{1,5}, Archana Amatya²

1 Department of Nursing, Dhaulagiri Prabhikhik Shikshya Pratishthan, Council for Technical Education and Vocational Training, Baglung, Nepal, **2** Central Department of Public Health, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal, **3** Department of Nutrition and Dietetics, Armed Police Force Hospital, Kathmandu, Nepal, **4** Center for Human Nutrition, Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, United States of America, **5** International Development Program, Nitze School of Advanced of International Studies (SAIS), Johns Hopkins University, Washington DC, United States of America

* shresthavintuna@gmail.com



Abstract

Background

Dietary diversity can play an important role in providing essential nutrients for both mother and fetus during pregnancy. This study aimed to assess the factors associated with dietary diversity during pregnancy in the western hill region of Nepal.

Methods

A cross-sectional study of 327 pregnant women was conducted in an urban municipality of Baglung district in the western hill region of Nepal. A semi-structured questionnaire was used to collect information on household demographic and socioeconomic status, food taboos, household food security status, nutrition-related knowledge in pregnancy, and women's empowerment. Women consuming ≥ 5 of 10 food groups in the past 24 hours were defined as consuming a diverse diet using the Minimum Dietary Diversity Score for Women (MDD-W) tool. Bivariate and multivariate logistic regression was used to estimate crude odds ratio (cOR) and adjusted odds ratios (aOR) and 95% confidence intervals (CIs) to understand factors associated with dietary diversity.

Results

Almost 45% (95% CI: 39.6–50.4) of the participants did not consume a diverse diet and the mean dietary diversity score was 4.76 ± 1.23 . Multivariable analysis revealed that women

OPEN ACCESS

Citation: Shrestha V, Paudel R, Sunuwar DR, Lyman AL, Manohar S, Amatya A (2021) Factors associated with dietary diversity among pregnant women in the western hill region of Nepal: A community based cross-sectional study. PLoS ONE 16(4): e0247085. <https://doi.org/10.1371/journal.pone.0247085>

Editor: Abid Hussain, International Centre for Integrated Mountain Development (ICIMOD), NEPAL

Received: September 21, 2020

Accepted: February 1, 2021

Published: April 8, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0247085>

Copyright: © 2021 Shrestha et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



FINDINGS




Fig 2. Conceptual framework on factors associated with dietary diversity.

PREDICTORS OF ADEQUATE DIETARY DIVERSITY (MULTIVARIABLE LOGISTIC REGRESSION ANALYSIS)

	aOR (95% C.I.)
Greater empowerment (vs. less)	4.3 (1.9, 9.9)
Wealth (upper vs. lower tertile)	5.1 (2.7, 9.3)
Greater nutrition knowledge (vs. less)	1.9 (1.1, 3.4)
Joint families (vs. nuclear)	2.7 (1.4, 5.1)



Study 3. Dietary diversity and nutrient adequacy among lactating women in peri-urban Nepal

- Objective: Assess the adequacy of the micronutrient intakes of lactating women in peri-urban Nepal
- N=500 randomly selected lactating women 17-44 years of age
- 3 x 24 hour recalls
- Mean probability of adequacy for 11 micronutrients was calculated

Public Health Nutrition: 18(17), 3201–3210

doi:10.1017/S136898001500067

Low dietary diversity and micronutrient adequacy among lactating women in a peri-urban area of Nepal

Sigrun Henjum^{1,*}, Liv Elin Torheim¹, Andrew L Thorne-Lyman^{2,3}, Ram Chandyo^{4,5}, Wafaie W Fawzi^{2,6}, Prakash S Shrestha⁵ and Tor A Strand^{4,7}

¹Oslo and Akershus University College of Applied Sciences, PO Box 4, St. Olavs Plass, 0130 Oslo, Norway; ²Department of Nutrition, Harvard School of Public Health, Boston, MA, USA; ³WorldFish, Penang, Malaysia; ⁴Centre for International Health, University of Bergen, Bergen, Norway; ⁵Department of Child Health, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal; ⁶Departments of Epidemiology and Global Health and Population, Harvard School of Public Health, Boston, MA, USA; ⁷Medical Services Division, Innlandet Hospital Trust Lillehammer, Norway

Submitted 23 June 2014; Final revision received 21 January 2015; Accepted 23 January 2015; First published online 31 March 2015

Abstract

Objective: The main objectives were to assess the adequacy of the micronutrient intakes of lactating women in a peri-urban area in Nepal and to describe the relationships between micronutrient intake adequacy, dietary diversity and sociodemographic variables.

Design: A cross-sectional survey was performed during 2008–2009. We used 24 h dietary recall to assess dietary intake on three non-consecutive days and calculated the probability of adequacy (PA) of the usual intake of eleven micronutrients and the overall mean probability of adequacy (MPA). A mean dietary diversity score (MDDS) was calculated of eight food groups averaged over 3 d. Multiple linear regression was used to identify the determinants of the MPA.

Setting: Bhaktapur municipality, Nepal.

Subjects: Lactating women (*n* 500), 17–44 years old, randomly selected.

Results: The mean usual energy intake was 8464 (sd 1305) kJ/d (2023 (sd 312) kcal/d), while the percentage of energy from protein, fat and carbohydrates was 11%, 13% and 76%, respectively. The mean usual micronutrient intakes were below the estimated average requirements for all micronutrients, with the exception of vitamin C and Zn. The MPA across eleven micronutrients was 0.19 (sd 0.16). The diet was found to be monotonous (MDDS was 3.9 (sd 1.0)) and rice contributed to about 60% of the energy intake. The multiple regression analyses showed that MPA was positively associated with energy intake, dietary diversity, women's educational level and socio-economic status, and was higher in the winter.

Conclusions: The low micronutrient intakes are probably explained by low dietary diversity and a low intake of micronutrient-rich foods.

Keyword
Dietary diversity
Micronutrient adequacy
Lactating women
Peri-urban Nepal





PREVALENCE OF ADEQUACY WAS VERY LOW

Table 3 Micronutrient requirements, usual micronutrient intakes*, prevalence of adequacy and MPA in lactating women (*n* 466) aged 17–44 years, Bhaktapur municipality, Nepal, January 2008–February 2009

	Requirements†		Usual intake according to BLUP‡			Prevalence of adequacy (%)	
	EAR	SD	Mean	SD	Median	Mean	SD
Thiamin (mg/d)	1.2	0.12	0.78	0.23	0.78	5	0.8
Riboflavin (mg/d)	1.3	0.13	0.62	0.18	0.59	1	0.3
Niacin (mg/d)	13.0	1.95	11.40	3.34	11.10	31	1.6
Vitamin B ₆ (mg/d)	1.7	0.17	1.45	0.37	1.44	26.7	1.6
Folate (µg/d)	450	45	163.8	70.0	147.9	0.3	0.2
Vitamin B ₁₂ (µg/d)	2.4	0.24	0.49	0.53	0.31	1	0.3
Vitamin C (mg/d)	55.0	5.5	60.0	27.6	46.3	36	2.0
Vitamin A (RE/d)	450	90	188.9	220.2	104.4	11	1.3
Ca (mg/d)	800	100	430.0	203.8	390.8	6.6	1.0
Fe (mg/d)§	23.4	7.02	17.4	8.6	15.2	28	1.4
Zn (mg/d)	7.0	0.88	7.80	1.60	7.63	66	1.6
MPA across 11 micronutrients						0.19	0.16
Median						0.15	

MPA, mean probability of adequacy; EAR, Estimated Average Requirement; RE, retinol equivalent.

*The mean is the overall mean of each individual's usual intake based on 3 d recall.

†The EAR are for lactating women; the EAR for vitamins A, C, B₆ and B₁₂, folate, thiamin, riboflavin, niacin and Fe were from WHO/FAO 2004⁽⁴⁾. For Ca, the Institute of Medicine EAR from 2011 was used⁽²⁴⁾. For Zn, the International Zinc Nutrition Consultative Group EAR for a mixed or refined diet was used assuming 44 % bioavailability⁽²⁵⁾. The SD of requirements was calculated using the CV of requirements and the EAR; the CV were 12.5 % for Zn⁽²⁵⁾, 20 % for vitamin A, 15 % for niacin, 10 % for vitamins C, B₆, B₁₂, thiamin, riboflavin, folate^(2,8) and Ca⁽²⁴⁾, and 30 % for Fe⁽²⁰⁾.

‡Best linear unbiased predictor⁽²¹⁾.

§Assuming 5 % bioavailability.

||The mean probability of adequacy is the average of the probability of adequacy of the eleven micronutrients.

CONCLUSIONS

- Mean usual micronutrient intakes were below estimated average requirements for all micronutrients except vitamin C and zinc
- Mean probability of adequacy across eleven micronutrients was 0.19 (SD 0.16)
- Rice contributed to 60% of energy intake
- Low dietary diversity explains the low micronutrient intakes, even in this peri-urban population outside the Kathmandu Valley

STUDY 4. IN THE SAME POPULATION, INTAKE OF FE SUPPLEMENTS IN PREGNANCY PREDICTED HB STATUS DURING LACTATION AND BODY FE FOR SOME...

	Multiple adjusted* β	95% C.I.	<i>P</i>
Model 1: Hemoglobin			
Intake of supplements in pregnancy	0.29	(0.04, 0.54)	0.03
Model 2. Body Fe (mg Fe/kg body weight)			
Dietary Fe (mg)	0.03	(0.014, 0.045)	<0.01
Interaction: time since birth*Fe in pregnancy Intake of Fe supplements in pregnancy	2.69	(1.54, 3.84)	<0.01
Time since birth <6 months	2.72	(1.79, 3.65)	<0.01
Time since birth \geq 6 months	0.02	(-0.67, 0.78)	0.93
Mother's age	0.21	(0.14, 0.28)	<0.01
Literacy	0.81	(0.25, 1.39)	<0.01
Land ownership	0.74	(0.19, 1.30)	<0.01

*Both models included mother and child age, parity, literacy

Interaction between time since birth (dichotomous, cut-off 6 months) and iron supplements in pregnancy for at least 6 months



Study 5. Vitamin D insufficiency among Nepali infants is low despite high prevalence among their mothers

- Objective: Describe the status and predictors of vitamin D status in healthy Nepali mother and infant pairs
- N=500 randomly selected mother and infant pairs
- Plasma 25(OH)D concentrations measured by LC-MS/MS
- Mean daily solar radiation over last 3 months
- Prevalence of insufficiency 25(OH)D, <50 nmol/L
 - Infants: 3.6%
 - Mothers: 59.8%

Low Prevalence of Vitamin D Insufficiency among Nepalese Infants Despite High Prevalence of Vitamin D Insufficiency among Their Mothers

Johanne Haugen ^{1,2,*}, Manjeswori Ulak ³, Ram K. Chandyo ³, Sigrun Henjum ⁴, Andrew L. Thorne-Lyman ^{5,6,7}, Per Magne Ueland ^{8,9}, Øivind Midttun ^{8,9}, Prakash S. Shrestha ³ and Tor A. Strand ^{1,2,10}

¹ Innlandet Hospital Trust, Lillehammer 2609, Norway; tors@me.com

² Centre for International Health, University of Bergen, Bergen 5007, Norway

³ Department of Child Health, Institute of Medicine, Tribhuvan University, Kathmandu 8212, Nepal; manjeswori@gmail.com (M.U.); ram.chandyo@cih.uib.no (R.K.C.); prakashsunder@hotmail.com (P.S.S.)

⁴ Department of Nursing and Health Promotion, Oslo and Akershus University College of Applied Sciences, Oslo 0130, Norway; sigrun.henjum@hioa.no

⁵ Johns Hopkins Center for Human Nutrition, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD 21205, USA; athorne1@jhu.edu

⁶ WorldFish, P.O. Box 500 GPO, Penang 10670, Malaysia

⁷ Departments of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA

⁸ Department of Clinical Science, University of Bergen, Bergen 5007, Norway; per.ueland@ikb.uib.no (P.M.U.); oivind.midttun@bevital.no (Ø.M.)

⁹ Bevital AS, Bergen 5021, Norway

¹⁰ Department of Sports Science, Inland Norway University of Applied Sciences, Lillehammer 2604, Norway

* Correspondence: johanne_haugen@hotmail.com; Tel.: +47-982-67-656

Received: 1 November 2016; Accepted: 7 December 2016; Published: 21 December 2016

Abstract: Background: Describing vitamin D status and its predictors in various populations is important in order to target public health measures. Objectives: To describe the status and predictors of vitamin D status in healthy Nepalese mothers and infants. Methods: 500 randomly selected Nepalese mother and infant pairs were included in a cross-sectional study. Plasma 25(OH)D concentrations were measured by LC-MS/MS and multiple linear regression analyses were used to identify predictors of vitamin D status. Results: Among the infants, the prevalence of vitamin D

PREDICTORS OF MATERNAL 25(OH)D (N=499)

		Crude β	95% CI	p	Adj. β	95% CI	p
BMI (kg/m ²)		0.3	(-0.1, 0.6)	0.109	0.6	(0.2, 1.0)	0.002
Global solar radiation (MJ/m ² /d)		2.6	(1.8, 3.4)	<0.001	2.6	(1.8, 3.4)	<0.001
Mother's age (y)		-0.6	(-0.9, -0.2)	0.002	-0.7	(-1.1, -0.4)	<0.001

Other variables tested in crude models but not significantly associated with 25(OH)D included blood pressure, CRP, maternal average energy-intake, parity, occupational status

PREDICTORS OF INFANT 25(OH)D

		Crude β	95% CI	p	Adj. β	95% CI	p
Age of child (months)		-2.0	(-2.6, -1.3)	<0.001	-1.5	(-2.1, 0.9)	<0.001
Mothers 25(OH)D (nmol/L)		0.5	0.4, 0.6	<0.001	0.5	(0.4, 0.6)	<0.001
Mothers BMI (kg/m ²)		0.9	0.3, 1.5	0.004	0.6	(0.02, 1.2)	0.041

Other variables tested in crude models but not significantly associated with 25(OH)D included sex, anthropometric z-scores, birth weight, CRP, duration of exclusive breastfeeding, maternal average energy-intake, global solar radiation, mother's occupational status

POSSIBLE EXPLANATIONS

- Breastmilk traditionally perceived as a poor source of vitamin D..but depends on woman's status!
- Short half life of 25(OH)D...Could maternal transfer of vitamin D3 to the infant and lower production of maternal 25(OH)D be an explanation?
- Fortified foods?
 - All infants were breastfed, most also received complementary foods..
 - All 70 exclusively breastfed infants were sufficient...
- Supplement use by women was rare...
- Might it be oil massage in the sun/dermal synthesis?



Photo by Bal Krishna Thapa/THT, The Himalayan

ACKNOWLEDGEMENTS

- Respondents
- **Funding: USAID** through Feed the Future Innovation Lab for Collaborative Research on Nutrition (Mission & DC)
- **Management Entity:** Tufts University, Friedman School of Food Policy & Nutrition
- **Collaborating institutions:** Institute of Medicine & NARC
- **Investigative Team:** Drs. Keith West, Rolf Klemm, Devendra Gauchan, Ramesh Adhikari, Andrew Thorne- Lyman, Patrick Webb, Shibani Ghosh, Swetha Manohar, Angela KC, Monica Pasqualino
- **PoSHAN-JHU Technical Team:** Abhigyna Bhattarai, Binod Shrestha, Shiva Bhandari, Sumanta Neupane, Ruchita Rajbhandary, Raman Shrestha, Hem Raj Paudel, Hari Krishna Shah, Dev Raj Gautam, Dev Mandal, Sudeep Shrestha, Chandni Karmacharya, NNIPS
- **Data Collection:** New Era Pvt. Ltd; Nepali Technical Assistance Group
- **Government of Nepal** for their support of this study

Work in Bhaktapur:

Dr. Tor Strand, U. Bergen

Dr. Mari Hysing, U. Bergen

Dr. Ingrid Kvested, Regional Center for Child and
Youth Mental Health and Child Welfare, NORCE
Norwegian Research Centre, Bergen Norway

Johanna Haugan, U. Bergen


Dr. Wafaie Fawzi, Harvard T.H. Chan School of
Public Health

Institute of Medicine at Tribhuvan University:

- Dr. Merina Shrestha
- Dr. Prakash Sundar Shrestha
- Dr. Laxman Shrestha
- Dr. Ram Chandyo
- Dr. Manjuswari Ulak
- Dr. Archana Amatya
- Dr. Rajan Paudel
- Vintuna Shrestha

Co-funding by the Bill & Melinda Gates Foundation, Sight & Life, Norwegian Council of Universities' Committee for Development Research and Education, Research Council of Norway, and GCRieber Funds, and National Institutes of Health (T32 #DK007703)

Overcoming Micronutrient Deficiencies in Women and Children



Omar Dary, USAID/Bureau for Global Health, Maternal and Child Health and Nutrition Office

May 19th, 2021

MOTHERS TRANSFER MICRONUTRIENTS DURING PREGNANCY AND LACTATION

Breast milk as food:

- **Macronutrients:** Protein, fat, carbohydrates (energy and building blocks)
- **Micronutrients:** vitamins and minerals (catalysis and functions)
- **Protective substances:** antibodies, non-digestible compounds that promote a healthy intestinal flora

	Nutrient	↑ in milk
Fortification vehicle		
Flour	B12	+++
Fish sauce	BI	+++
Sugar	A	+++
Salt	I	+++
Food		
Red palm oil	A	+++
B-carotene	A	+/-
Fish	DHA	+
Supplements		
LNS	Multiple	++
B12	B12	+


Content does not change with intake for: Folate
Iron, copper, zinc, and calcium

Source: Lindsay Allen, USDA and University of California at Davis






PHYSICAL GROWTH MIGHT BE DETERMINED AT THE MOMENT OF BIRTH



Child stunting evolution in a poor rural community of Guatemala where mothers (pregnancy) and children (starting at 6 months) received CSB, LNS, or MNP's.

Source: Olney *et al.* *J Nutr* 2018; 148:1493.


INTELLECTUAL DISABILITY DUE TO IODINE DEFICIENCY HAS DISAPPEARED



Source: Jonathan Gorstein, Iodine Global Network.




or dairy products from animals with good iodine status







REDUCTION OF XEROPHTHALMIA (VIT. A DEFICIENCY) IN GUATEMALA



SUN LIGHT OR GOOD STATUS DURING PREGNANCY IS NEEDED FOR VIT. D




Source: Choubey et al., *Int J Dent Case Reports* 2013; 3(2):16-19



Source: Anastassios Pittas, Tuft's University. Conference about impact of vitamin D supplementation and prevention of diabetes



NEURAL TUBE DEFECTS ARE PREVENTED BEFORE PREGNANCY



Source: Jorge Rosenthal, CDC


Neural tube defects, the most common birth defect. It appears within 28 days after conception, consequence of genetic vulnerability, and folate and vit. B₁₂ deficiencies, among others.

Incapacities:

- Leg paralysis
- Hydrocephaly
- Bad control of bladder and intestinal evacuations
- Learning difficulties




Birth defects in Chile after introduction of wheat flour fortified with folic acid






IRON SUPPLEMENTATION IN YOUNG CHILDREN

Excessive iron supply favors pathogenic intestinal bacteria in Kenyan children



Source: Jaeggi *et al.*,
doi:10.1136/gutjnl-2014-
307720

GOOD NUTRITION IS A GOAL ALONG THE LIFE COURSE FOR THE FAMILY



Source: USAID/BFS, adapted from ACC/SCN (2000) Fourth Report on the World Nutrition Situation. Geneva: ACC/SCN in collaboration with the International Food Policy Research Institute.

CONCLUSIONS

1. The nutritional and health status of women before and during pregnancy is needed for their well-being and survival, as well as the physical, mental, and social development of their children.
2. Micronutrients (vitamins and minerals) and other essential nutrients are transferred from mothers to children during pregnancy and through breast milk.
3. Intake of micronutrients from mothers is reflected in the breast milk content, except for calcium (this comes from the mother's bones), folate, iron, copper, zinc, and therefore they must be transferred during pregnancy.
4. Micronutrient interventions, mostly food fortification, have been successful to improve the women's micronutrient status, and therefore the status of the offspring.
5. Trying to improve the nutritional status during pregnancy may be late for some micronutrients, Therefore, attention to the family's diet and habits is important.



FEED^{THE}FUTURE
The U.S. Government's Global Hunger & Food Security Initiative

Q&A

THANK YOU

- To register for upcoming webinars, you can visit **NutritionInnovationLab.org** or **AdvancingNutrition.org**. More details coming soon!
- Recordings and slides for each webinar will also be posted on our websites.



FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative

www.feedthefuture.gov



USAID
FROM THE AMERICAN PEOPLE