

Definitions:

STUNTING:

Proportion of children with a length-/height-for-age z-score more than 2 standard deviations below the mean.

UNDERWEIGHT:

Proportion of children with a weight-for-age z-score more than 2 standard deviations below the mean.

MODERATE ACUTE MALNUTRITION:

Proportion of children with a weight-for-length/height z-score between 2 and 3 standard deviations below the mean.

WEALTH:

A wealth index was constructed using the DHS method, including data on a household's ownership of selected assets; materials used for housing construction; and types of water access and sanitation facilities. Wealth quintiles are reported here.

FEMALE CAREGIVER'S EDUCATION:


Categories are defined using the highest grade of schooling completed by the female caregiver. Primary 1 incomplete includes completing grades 1, 2, 3, or 4; primary 1 complete includes completing grades 5, 6, or 7; and any secondary or higher includes completing grades 8, 9, 10, 11, 12, or any higher education.

This infographic was produced as a part of a study that was conducted by the Friedman School of Nutrition Science and Policy at Tufts University, in collaboration with University Lúrio, Mozambican National Institute for Health, and the Association for Food and Nutrition Security (ANSA), under the Feed the Future Innovation Lab for Nutrition.

The study was conducted in 10 districts of Nampula province where USAID supports programs in nutrition and agriculture.

Support for this publication was provided by the Feed the Future Innovation Lab for Nutrition, which is funded by the United States Agency for International Development, CSI and USAID Mission under grant ID:AID-OAA-L-10-00006.

nutritioninnovationlab.org



ANTHROPOMETRY
of Children 6-59 months

|||||


STUNTING





AGE



WEALTH



Female Caregiver's EDUCATION




UNDERWEIGHT




AGE



WEALTH



Female Caregiver's EDUCATION



Moderate Acute MALNUTRITION

CHILDREN

2%

of children had moderate acute malnutrition. Children with severe acute malnutrition were excluded from enrollment.